

A topological theory of Maslov indices for Lagrangian and symplectic paths

M. de Gosson and S. de Gosson
maurice.degossion@gmail.com

Abstract

We propose a topological theory of the Maslov index for lagrangian and symplectic paths based on a minimal system of axioms. We recover, as particular cases, the Hörmander and the Robbin–Salomon indices.

1 Introduction

The theory of the Maslov index has recently found new applications in intersection theory.

1.1 General notations

We essentially use the notations of Leray [18]. Let $X = \mathbb{R}^n$; the product $X \times X^*$ is endowed with the canonical symplectic structure defined by:

$$\omega(z, z') = \langle p, x' \rangle - \langle p', x \rangle$$

if $z = (x, p)$, $z' = (x', p')$. The symplectic group of $(X \times X^*, \omega)$ will be denoted by $Sp(n)$: we have $s \in Sp(n)$ if and only if $\omega(sz, sz') = \omega(z, z')$ for all z, z' . The Poincaré group $\pi_1(Sp(n))$ is isomorphic to $(\mathbb{Z}, +)$ and $Sp(n)$ has thus covering groups $Sp_q(n)$ of all orders $q = 1, 2, \dots, \infty$ ($Sp_\infty(n)$ is the universal covering group of $Sp(n)$). The unitary group $U(n)$ is identified with a subgroup of $Sp(n)$ by defining the action of $u = A + iB$ on $z = x + p$ by the formula

$$\begin{pmatrix} x' \\ p' \end{pmatrix} = \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}$$

We denote by $\Lambda(n)$ the Lagrangian Grassmannian of $(X \times X^*, \omega)$: $\ell \in \Lambda(n)$ if and only if ℓ is an n dimensional subspace of $X \times X^*$ on which the

form ω vanishes. The manifold $\Lambda(n)$ is connected and compact manifold and has dimension $n(n+1)/2$. Its Poincaré group $\pi_1(\Lambda(n))$ is isomorphic to $(\mathbb{Z}, +)$ and $\Lambda(n)$ thus has coverings $\Lambda_q(n)$ of all orders q . Its universal covering $\Lambda_\infty(n)$ is sometimes called the “Maslov bundle”.

The group $Sp(n)$, and hence $U(n)$, acts on $\Lambda(n)$. The action of $U(n)$, and hence that of $Sp(n)$, is transitive: the orbit of every $\ell \in \Lambda(n)$ under the action of $Sp(n)$ is $\Lambda(n)$.

We shall denote by $St_\ell(n)$ the stabilizer group of a Lagrangian ℓ in $Sp(n)$:

$$St_\ell(n) = \{s \in Sp(n) : s\ell = \ell\}.$$

One can show (see [1, 5, 18]) that the fibration $Sp(n)/St_{X^*}(n) = \Lambda(n)$ defines an isomorphism

$$\mathbb{Z} \cong \pi_1[Sp(n)] \longrightarrow \pi_1[\Lambda(n)] \cong \mathbb{Z}$$

which is multiplication by 2 on $\mathbb{Z}$. It follows (see [18], Theorem 3,3^o, p.36) that the action of $Sp(n)$ on $\Lambda(n)$ can be lifted to a transitive action of the universal covering $Sp_\infty(n)$ on the Maslov bundle $\Lambda_\infty(n)$ such that

$$(\alpha s_\infty)\ell_\infty = \beta^2(s_\infty\ell_\infty) = s_\infty(\beta^2\ell_\infty) \quad (1)$$

for all $(s_\infty, \ell_\infty) \in Sp_\infty(n) \times \Lambda_\infty(n)$; α (resp. β) is the generator of $\pi_1(Sp(n))$ (resp. $\pi_1(\Lambda(n))$) whose image in $\mathbb{Z}$ is $+1$.

The Lagrangian Grassmannian $\Lambda(n)$ is a stratified manifold; for $\ell \in \Lambda$ and k an integer, $0 \leq k \leq n$,

$$\Lambda_\ell(n, k) = \{\ell' \in \Lambda(n) : \dim(\ell \cap \ell') = k\}$$

is the stratum of $\Lambda(n)$ of order k , relative to the Lagrangian plane ℓ . We will also use the following notations:

$$\Lambda^2(n; k) = \bigcup_{\ell \in \Lambda} \Lambda_\ell(n, k) \quad , \quad \Lambda_\infty^2(n; k) = \pi^{-1}(\Lambda^2(n; k))$$

that is:

$$\begin{aligned} \Lambda^2(n; k) &= \{(\ell, \ell') \in \Lambda(n)^2 : \dim(\ell \cap \ell') = k\} \\ \Lambda_\infty^2(n; k) &= \{(\ell_\infty, \ell'_\infty) \in \Lambda_\infty(n)^2 : \dim(\ell \cap \ell') = k\}. \end{aligned}$$

In particular, $\Lambda_\infty^2(n; 0)$ is the set of pairs $(\ell_\infty, \ell'_\infty)$ of elements of $\Lambda_\infty(n)$ whose projections ℓ and ℓ' are transversal. One proves (see for instance [26]) that $\Lambda_\ell(n; 0)$ is an open subset of $\Lambda(n)$ and that the $\Lambda_\ell(n; k)$ are, for

$0 \leq k \leq n$, connected submanifolds of $\Lambda(n)$, of codimension $k(k+1)/2$. The closed set

$$\Sigma_\ell = \Lambda(n) \setminus \Lambda_\ell(n; 0) = \bigcup_{k=1}^n \Lambda_\ell(k)$$

is called the “Maslov cycle relative to ℓ ”. It is the set of Lagrangians that are not transverse to ℓ . When $\ell = X^*$ we call Σ_ℓ the *Maslov cycle*, and denote it by Σ .

1.2 Cohomological notations

It will be economical to use the following elementary notations from the singular Alexander–Spanier cohomology: let E be a set, k a positive integer or 0, and $(G, +)$ an abelian group. We shall call the functions $f : E^{k+1} \rightarrow G$ *k-cochains on E with values in G* (or simply *k-cochains*, or *cochains*). The coboundary ∂f of a k -cochain is the $(k+1)$ -cochain defined by:

$$\partial f(a_0, \dots, a_{k+1}) = \sum_{j=0}^{k+1} (-1)^j f(a_0, \dots, \widehat{a_j}, \dots, a_{k+1}),$$

where the cap $\widehat{}$ deletes the term it covers. We evidently have that $\partial^2 f = 0$ for every cochain f . A k -cochain f is a *coboundary* if there exists a cochain g such that $f = \partial g$; a cochain f is a *cocycle* if $\partial f = 0$. We thus have that every coboundary is a cocycle.

2 The Keller-Maslov index

In this section we recall the definition of the Keller-Maslov index (see [1, 18, 24] for a thorough study).

Using the ideas of Keller [16], Maslov introduced in [21] an index, whose definition was clarified by [1] and extended by Leray [17, 18] and one of the authors [7]. (Dazord [5] has given similar results in the more general case of symplectic bundles.) The Keller-Maslov index is designed to count the number of intersections of a Lagrangian loop γ with the Maslov cycle Σ . It is defined in the following way. Let $W(n, \mathbb{C})$ be the submanifold of $U(n, \mathbb{C})$ consisting of symmetric matrices:

$$W(n, \mathbb{C}) = \{u \in U(n, \mathbb{C}) : u = u^t\}.$$

($u^t = (u^*)^{-1}$ the transpose of u). The mapping

$$\Lambda(n) \ni \ell = uX^* \mapsto uu^t \in U(n, \mathbb{C})$$

is a homeomorphism

$$\Lambda(n) \ni \ell = uX^* \longmapsto uu^t \in W(n, \mathbb{C})$$

The composition of the isomorphism $\pi_1(\Lambda(n)) \cong \pi_1(W(n, \mathbb{C}))$ induced by this homeomorphism with the isomorphism

$$\pi_1(W(n, \mathbb{C})) \ni [\gamma] \longmapsto \frac{1}{2\pi i} \oint_{\gamma} \frac{d(\det w)}{\det w} \in \mathbb{Z} \quad (2)$$

is an isomorphism

$$\text{ind} : \pi_1(\Lambda(n)) \ni [\gamma] \xrightarrow{\cong} \text{ind}[\gamma] \in (\mathbb{Z}, +).$$

By definition, The Keller-Maslov index on $\Lambda(n)$ is the mapping m that to each Lagrangian loop γ associate the integer:

$$m(\gamma) = \text{ind}[\gamma] = \frac{1}{2\pi i} \oint_{\gamma} \frac{d(\det w)}{\det w}. \quad (3)$$

We can easily show that $m(\gamma)$ is zero in the strata of $\Lambda(n)$:

$$\text{Im } \gamma \subset \Lambda_{\ell}(n; k) \implies m(\gamma) = 0.$$

Let β be the generator of $\pi_1(\Lambda(n))$ whose natural image in $\mathbb{Z}$ is $+1$. Its Maslov index is

$$m(\beta) = 1. \quad (4)$$

The Keller-Maslov index is an homotopic invariant and in particular $m(\gamma) = 0$ if γ is contractible to a point. Furthermore, this index possesses the following additivity property:

$$m(\gamma * \gamma') = m(\gamma) + m(\gamma') \quad (5)$$

for all consecutive loops γ and γ' ($\gamma * \gamma'$ being the loop γ followed by the loop γ').

One should note the difference between the Keller-Maslov index m on $\Lambda(n)$ and the index m_V of loop on a Lagrangian submanifold V in $X \times X^*$. The index m_V is defined in the following way: Let $\ell(\cdot)$ be the continuous mapping that to each $z \in V$ associates the tangent space at z : $\ell(z) = T_z V$. This mapping associated to each loop γ_V in V a loop γ in $\Lambda(n)$, and by definition:

$$m_V(\gamma_V) = m(\gamma). \quad (6)$$

Souriau proved in [25] that

$$m_V(\gamma_V) \text{ is an even integer when the manifold } V \text{ is oriented.} \quad (7)$$

In fact, we have the more precise result (see de Gosson [8]). In what follows we denote by $\Lambda_{2q}(n)$ the connected $2q$ -fold covering of $\Lambda(n)$, $q = 1, 2, \dots$.

Proposition 1 *Suppose that the mapping $\ell(\cdot) : V \longrightarrow \Lambda(n)$ is lifted to a continuous mapping $\ell_{2q}(\cdot) : V \longrightarrow \Lambda_{2q}(n)$. Then*

$$m_V(\gamma_V) \equiv 0 \pmod{2q}$$

for every loop γ_V in V .

Note that if $\ell(\cdot)$ is lifted to a continuous mapping $\ell_\infty(\cdot) : V \longrightarrow \Lambda_\infty(n)$ we have $m(\gamma_V) = 0$ for every loop γ_V in V since the universal covering $\Lambda_\infty(n)$ is simply connected.

3 Definition of the indices μ_Λ and μ_{Sp}

Let $\mathcal{C}(\Lambda(n))$ be the set of continuous paths $[0, 1] \longrightarrow \Lambda(n)$. If λ and λ' are two consecutive paths (i.e., if $\lambda(1) = \lambda'(0)$) we shall denote $\lambda * \lambda'$ the path λ followed by the path λ' :

$$\lambda * \lambda'(t) = \begin{cases} \lambda(2t) & \text{if } 0 \leq t \leq \frac{1}{2} \\ \lambda'(2t - 1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases}$$

We denote by λ^{-1} the inverse of the path λ : $\lambda^{-1}(t) = \lambda(1 - t)$. Finally, we shall write $\lambda \sim \lambda'$ when the paths λ and λ' are homotopic with fixed endpoints.

3.1 Axioms for the indices μ_Λ

Definition 2 *A "Lagrangian intersection index" is a mapping*

$$\mu_\Lambda : \mathcal{C}(\Lambda(n)) \times \Lambda(n) \ni (\lambda, \ell) \longmapsto \mu_{\Lambda(n)}(\lambda, \ell) \in \mathbb{Z}$$

having the following four properties:

(L₁) homotopic invariance: *If the paths λ and λ' in $\Lambda(n)$ have the same endpoints, then $\mu_\Lambda(\lambda, \ell) = \mu_\Lambda(\lambda', \ell)$ if and only if $\lambda \sim \lambda'$;*

(L₂) *Additivity under composition: If λ and λ' are two consecutive paths, then for all $\ell \in \Lambda(n)$:*

$$\mu_\Lambda(\lambda * \lambda', \ell) = \mu_\Lambda(\lambda, \ell) + \mu_\Lambda(\lambda', \ell)$$

(L₃) *Zero in the stratum. If the path λ remains in the same stratum $\Lambda_\ell(n; k)$, then $\mu_\Lambda(\lambda, \ell)$ is zero, i.e.,*

$$\dim \lambda(t) \cap \ell = k \quad (0 \leq t \leq 1) \implies \mu_\Lambda(\lambda, \ell) = 0.$$

(L₄) *Restriction to loops. If γ is a loop, then $\mu_\Lambda(\gamma, \ell) = 2m(\gamma)$ ($m(\gamma)$ the Maslov index of γ) for all ℓ .*

(We shall see in 6.1 that the condition $\mu_\Lambda(\gamma, \ell) = 2m(\gamma)$, and not $\mu_\Lambda(\gamma, \ell) = m(\gamma)$, is necessary for an index satisfying the axioms (L₁–L₃) to be an integer)

We note in particular that the axioms (L₂) and (L₄) imply that the index is antisymmetric, i.e. ,

$$\mu_\Lambda(\lambda^{-1}, \ell) = -\mu_\Lambda(\lambda, \ell). \quad (8)$$

Indeed, by (L₂) we have

$$\mu_{\Lambda(n)}(\lambda * \lambda^{-1}, \ell) = \mu_{\Lambda(n)}(\lambda, \ell) + \mu_{\Lambda(n)}(\lambda^{-1}, \ell)$$

but since the loop $\lambda * \lambda^{-1} = \gamma$ is homotopic to a point the axiom (L₄) implies that

$$\mu_{\Lambda(n)}(\lambda * \lambda^{-1}, \ell) = 2m(\gamma) = 0.$$

The system of axioms (L₁)–(L₄) is in fact equivalent to the system of axioms obtained by replacing (L₁) by an apparently stronger condition (L'₁) we are going to describe below. Let us first define the notion of “homotopy in strata”:

Definition 3 *Two Lagrangian paths λ and λ' are said to be “homotopic in the strata relative to ℓ ” (denoted $\lambda \approx_\ell \lambda'$) if there exists a continuous mapping $h : [0, 1] \times [0, 1] \longrightarrow \Lambda(n)$ such that*

$$h(t, 0) = \lambda(t) \quad , \quad h(t, 1) = \lambda'(t) \quad \text{for } 0 \leq t \leq 1$$

and two integers k_0, k_1 ($0 \leq k_0, k_1 \leq n$) such that

$$h(0, s) \in \Lambda_\ell(n; k_0) \quad \text{and} \quad h(1, s) \in \Lambda_\ell(n; k_1) \quad \text{for } 0 \leq s \leq 1.$$

The intersection indices μ_Λ have the following property that makes (L₁) more precise:

Proposition 4

(L'₁) *If the paths λ and λ' are homotopic in strata relative to ℓ , then $\mu_\Lambda(\lambda, \ell) = \mu_\Lambda(\lambda', \ell)$, i.e. ,*

$$\lambda \approx_\ell \lambda' \implies \mu_\Lambda(\lambda, \ell) = \mu_\Lambda(\lambda', \ell). \quad (9)$$

Proof. Suppose that $\lambda \approx_\ell \lambda'$ and define the paths ε_0 and ε_1 joining $\lambda'(0)$ to $\lambda(0)$ and $\lambda(1)$ to $\lambda'(1)$, respectively, by $\varepsilon_0(s) = h(0, 1-s)$ and $\varepsilon_1(s) = h(1, s)$ ($0 \leq s \leq 1$). Then $\lambda * \varepsilon_1 * \lambda'^{-1} * \varepsilon_0$ is homotopic to a point, and hence, in view of (L₂) and (L₄):

$$\mu_\Lambda(\lambda, \ell) + \mu_\Lambda(\varepsilon_1, \ell) + \mu_\Lambda(\lambda'^{-1}, \ell) + \mu_\Lambda(\varepsilon_0, \ell) = 0.$$

But, in view of (L₃)

$$\mu_\Lambda(\varepsilon_1, \ell) = \mu_\Lambda(\varepsilon_0, \ell) = 0$$

and thus

$$\mu_\Lambda(\lambda, \ell) + \mu_\Lambda(\lambda'^{-1}, \ell) = 0$$

hence $\mu_\Lambda(\lambda, \ell) = \mu_\Lambda(\lambda', \ell)$ in view of (8). ■

3.2 A property of relative uniqueness

The axioms (L₁–L₄) do not guaranty the uniqueness of an intersection index. Indeed, if μ_Λ is such an index, then the function $\mu'_{\Lambda(n)} : \mathcal{C}(\Lambda(n)) \times \Lambda(n) \longrightarrow \mathbb{Z}$ defined by

$$\mu'_\Lambda(\lambda, \ell) = \mu(\lambda, \ell) + \dim(\lambda(1) \cap \ell) - \dim(\lambda(0) \cap \ell)$$

is also an intersection index. We will prove in this section a result of “relative uniqueness” modulo a 1-cochain.

Let us introduce the following notations: for every pair of points (ℓ_i, ℓ_j) of $\Lambda(n)$ we denote λ_{ij} or λ'_{ij} an arbitrary element of $\mathcal{C}(\Lambda(n))$ joining ℓ_i to ℓ_j . The opposite path λ_{ij}^{-1} of λ_{ij} will then be denoted λ_{ji} . Using these notations we can formulate a theorem relating all intersection indices to one another. That is, show that there exists some function allowing us to relate one intersection index to another, thereby giving us a relative uniqueness theorem for intersection indices.

Theorem 5 *Let μ_Λ and μ'_Λ be two intersection indices on $\Lambda(n)$. For every $\ell \in \Lambda(n)$ there exists a function $f : \mathbb{N} \longrightarrow \mathbb{Z}$ such that:*

$$\mu_\Lambda(\lambda_{01}, \ell) - \mu'_\Lambda(\lambda_{01}, \ell) = f(\dim(\ell_0 \cap \ell)) - f(\dim(\ell_1 \cap \ell)). \quad (10)$$

To prove this fact we have to make sure that the left-hand side of (10) does not depend on anything else than ℓ_0, ℓ_1 (and of course ℓ). We have the essential

Lemma 6 *for ℓ fixed, the difference $\mu_\Lambda(\lambda_{01}, \ell) - \mu'_\Lambda(\lambda_{01}, \ell)$ depends only on (ℓ_0, ℓ_1) , and the 1-cochain δ_ℓ on $\Lambda(n)$ given by:*

$$\delta_\ell(\ell_0, \ell_1) = \mu_\Lambda(\lambda_{01}, \ell) - \mu'_\Lambda(\lambda_{01}, \ell) \quad (11)$$

is an antisymmetric cocycle:

$$\partial\delta_\ell(\ell_0, \ell_1, \ell_2) = 0 \quad \text{and} \quad \delta_\ell(\ell_0, \ell_1) = -\delta_\ell(\ell_1, \ell_0) \quad (12)$$

for all $(\ell_0, \ell_1, \ell_2) \in \Lambda(n)^3$.

Proof of the lemma. Suppose that we replace the path λ_{01} with another path λ'_{01} joining ℓ_0 to ℓ_1 . Then, there exists a loop γ_1 passing by ℓ_1 and such that $\lambda'_{01} \sim \lambda_{01} * \gamma_1$ and we have, by the axioms (L₁) and (L₄):

$$\begin{cases} \mu_\Lambda(\lambda'_{01}, \ell) = \mu_\Lambda(\lambda_{01}, \ell) + 2m(\gamma_1) \\ \mu'_\Lambda(\lambda'_{01}, \ell) = \mu'_\Lambda(\lambda_{01}, \ell) + 2m(\gamma_1) \end{cases}.$$

Hence, the difference

$$\mu_\Lambda(\lambda'_{01}, \ell) - \mu'_\Lambda(\lambda'_{01}, \ell) = \mu_\Lambda(\lambda_{01}, \ell) - \mu'_\Lambda(\lambda_{01}, \ell) = \delta_\ell(\ell_0, \ell_1)$$

depends only on ℓ_0 and ℓ_1 (and ℓ). We now show that the coboundary of that 1-cochain is zero: $\partial\delta_\ell = 0$. Let (ℓ_0, ℓ_1, ℓ_2) be an arbitrary triple of points in $\Lambda(n)$. By the properties (L₂) and (L₄) we see that

$$\begin{cases} \mu_\Lambda(\lambda_{01}, \ell) - \mu_\Lambda(\lambda_{02}, \ell) + \mu_\Lambda(\lambda_{12}, \ell) = 2m(\gamma_0) \\ \mu'_\Lambda(\lambda_{01}, \ell) - \mu'_\Lambda(\lambda_{02}, \ell) + \mu'_\Lambda(\lambda_{12}, \ell) = 2m(\gamma_0) \end{cases} \quad (13)$$

where $\gamma_0 = \lambda_{01} * \lambda_{12} * \lambda_{20}$ is a loop with origin ℓ_0 . Subtracting the second equality (13) from the first and using the definition of δ_ℓ we obtain:

$$\partial\delta_\ell(\ell_0, \ell_1, \ell_2) = 0.$$

The antisymmetry of δ_ℓ is an immediate consequence of (8). ■

Proof of the theorem. (1) By the definition of δ_ℓ we have

$$\mu_\Lambda(\lambda_{01}, \ell) - \mu_\Lambda(\lambda'_{01}, \ell) = \delta_\ell(\ell_0, \ell_1).$$

Since δ_ℓ is a cocycle this equality can be written as

$$\mu_\Lambda(\lambda_{01}, \ell) - \mu_\Lambda(\lambda'_{01}, \ell) = \delta_\ell(\ell_0, \ell) - \delta_\ell(\ell_1, \ell).$$

We shall show that the 0-cochain $\ell_0 \mapsto \delta_\ell(\ell_0, \ell)$ is constant in each stratum $\Lambda_\ell(n; k)$ and the theorem will follow. We see, by the definition of δ_ℓ that

$$\delta_\ell(\ell_0, \ell) = \mu_\Lambda(\lambda, \ell) - \mu'_\Lambda(\lambda, \ell)$$

where λ is an arbitrary path joining ℓ_0 to ℓ in $\Lambda(n)$. Let λ' be a path joining ℓ_0 to ℓ'_0 in $\Lambda_\ell(n; k)$. Then

$$\delta_\ell(\ell'_0, \ell) = \mu_\Lambda(\lambda' * \lambda, \ell) - \mu'_\Lambda(\lambda' * \lambda, \ell)$$

and in view of the additivity property (L₂)

$$\delta_\ell(\ell'_0, \ell) = \mu_\Lambda(\lambda', \ell) - \mu'_\Lambda(\lambda', \ell) + \delta_\ell(\ell_0, \ell).$$

But, by the axiom (L₃)

$$\mu_\Lambda(\lambda', \ell) = \mu'_\Lambda(\lambda', \ell) = 0$$

and we thus have $\delta_\ell(\ell'_0, \ell) = \delta_\ell(\ell_0, \ell)$, from which we conclude that the mapping $\ell_0 \mapsto \delta_\ell(\ell_0, \ell)$ is constant in the stratum $\Lambda_\ell(n; k)$. ■

3.3 The signature cocycle

We now present an interesting result that will generalize the axiom (L₁) and that will enable us to give a notion of signature for every triple of elements of elements in $\Lambda(n)$.

Lemma 7 *Let ℓ_0, ℓ_1, ℓ_2 be three elements of $\Lambda(n)$. Suppose that the paths λ_{ij} and λ'_{ij} ($0 \leq i, j \leq 2$) are such that*

$$\lambda_{01} * \lambda_{12} * \lambda_{20} \sim \lambda'_{01} * \lambda'_{12} * \lambda'_{20}. \quad (14)$$

Then both sums

$$\Sigma = \mu_\Lambda(\lambda_{01}, \ell_2) + \mu_\Lambda(\lambda_{12}, \ell_0) + \mu_\Lambda(\lambda_{20}, \ell_1)$$

and

$$\Sigma' = \mu_\Lambda(\lambda'_{01}, \ell_2) + \mu_\Lambda(\lambda'_{12}, \ell_0) + \mu_\Lambda(\lambda'_{20}, \ell_1)$$

are equal: $\Sigma = \Sigma'$. Furthermore, Σ does not depend on the choice of μ_Λ .

Proof. There exist loops γ_0^1 , γ_0^2 and γ_0^3 passing through ℓ_0 such that

$$\begin{cases} \lambda'_{01} \sim \gamma_0^1 * \lambda_{01} \\ \lambda'_{12} \sim \lambda_{10} * \gamma_0^2 * \lambda_{01} * \lambda_{12} \\ \lambda'_{20} \sim \lambda_{20} * \gamma_0^3 \end{cases}.$$

In view of (14) and the properties (L₁) and (L₄) we obtain

$$\begin{aligned} \Sigma' &= \Sigma + 2(m(\gamma_0^1) + m(\gamma_0^2) + m(\gamma_0^3)) \\ &= \Sigma + 2m(\gamma_0^1 * \gamma_0^2 * \gamma_0^3). \end{aligned}$$

But (14) implies that $\gamma_0^1 * \gamma_0^2 * \gamma_0^3 \sim 0$ and hence that $\mu_{\Lambda(n)}(\gamma_0^1 * \gamma_0^2 * \gamma_0^3) = 0$ and $\Sigma' = \Sigma$. It remains to prove that Σ is independent of the choice of index $\mu_{\Lambda(n)}$. If $\mu'_{\Lambda(n)}$ is another intersection index and Σ' the associated sum the relative uniqueness theorem 5 tells us that

$$\begin{aligned} \Sigma - \Sigma' &= f(\dim(\ell_0 \cap \ell_2)) - f(\dim(\ell_1 \cap \ell_2)) + f(\dim(\ell_1 \cap \ell_0)) \\ &\quad - f(\dim(\ell_2 \cap \ell_0)) + f(\dim(\ell_2 \cap \ell_1)) - f(\dim(\ell_0 \cap \ell_1)) \end{aligned}$$

that is: $\Sigma - \Sigma' = 0$. ■

This result motivates the following definition

Definition 8 Let $(\ell_0, \ell_1, \ell_2) \in \Lambda(n)^3$ and λ_{01} , λ_{12} , λ_{20} be elements of $\mathcal{C}(\Lambda(n))$ such that $\lambda_{01} * \lambda_{12} * \lambda_{20}$ be homotopic to a point. The sum

$$\text{sign}(\ell_0, \ell_1, \ell_2) = \mu_{\Lambda}(\lambda_{01}, \ell_2) + \mu_{\Lambda}(\lambda_{12}, \ell_0) + \mu_{\Lambda}(\lambda_{20}, \ell_1) \quad (15)$$

is called the signature of the triple (ℓ_0, ℓ_1, ℓ_2) . This signature is independent of the choice of intersection index μ_{Λ} on $\Lambda(n)$.

Remark 9 In Section 4.2 we will show that sign is just the Demazure–Kashiwara signature τ ([19, 20]) arising in symplectic geometry.

By (8), the signature is an antisymmetric 2-cochain

$$\text{sign}(\ell_{\varepsilon(0)}, \ell_{\varepsilon(1)}, \ell_{\varepsilon(2)}) = (-1)^{\text{sgn}(\varepsilon)} \text{sign}(\ell_0, \ell_1, \ell_2)$$

for every permutation ε of $\{0, 1, 2\}$. Furthermore, it possesses the following essential property

Proposition 10 *The signature is a 2-cocycle:*

$$\partial \text{sign}(\ell_0, \ell_1, \ell_2, \ell_3) = 0 \quad (16)$$

that is

$$\begin{aligned} \text{sign}(\ell_1, \ell_2, \ell_3) - \text{sign}(\ell_0, \ell_2, \ell_3) \\ + \text{sign}(\ell_0, \ell_1, \ell_3) - \text{sign}(\ell_0, \ell_1, \ell_2) = 0. \end{aligned}$$

Proof. By the definition of the coboundary operator ∂ we have

$$\begin{aligned} \partial \text{sign}(\ell_0, \ell_1, \ell_2, \ell_3) &= \text{sign}(\ell_1, \ell_2, \ell_3) - \text{sign}(\ell_0, \ell_2, \ell_3) \\ &\quad + \text{sign}(\ell_0, \ell_1, \ell_3) - \text{sign}(\ell_0, \ell_1, \ell_2) \end{aligned}$$

and a calculation making use of (8) and the axiom (L₂) gives us

$$\begin{aligned} \partial \text{sign}(\ell_0, \ell_1, \ell_2, \ell_3) &= \\ &\mu_{\Lambda(n)}(\gamma_1, \ell_3) + \mu_{\Lambda(n)}(\gamma_2, \ell_1) + \mu_{\Lambda(n)}(\gamma_3, \ell_2) + \mu_{\Lambda(n)}(\gamma_4, \ell_0) \end{aligned}$$

where $\gamma_1 = \gamma_{12} * \gamma_{20} * \gamma_{01}$, $\gamma_2 = \gamma_{23} * \gamma_{30} * \gamma_{02}$, $\gamma_3 = \gamma_{31} * \gamma_{10} * \gamma_{03}$ and $\gamma_4 = \gamma_{32} * \gamma_{21} * \gamma_{13}$ are loop, hence by (L₄)

$$\partial \text{sign}(\ell_0, \ell_1, \ell_2, \ell_3) = 2(m(\gamma_1) + m(\gamma_2) + m(\gamma_3) + m(\gamma_4)).$$

The loops $\gamma_1, \gamma_2, \gamma_3$ and γ_4 being contractible to a point, the Maslov indices of this loops are zero and thereby $\partial \text{sign} = 0$. ■

3.4 The indices μ_{Sp}

We shall denote by $\mathcal{C}(Sp(n))$ the set of continuous paths $[0, 1] \longrightarrow Sp(n)$.

Definition 11 *A symplectic intersection index is a mapping*

$$\mu_{Sp} : \mathcal{C}(Sp(n)) \times \Lambda(n) \ni (\sigma, \ell) \longmapsto \mu_{Sp}(\sigma, \ell) \in \mathbb{Z}$$

satisfying the following four axioms:

- (S₁) *Homotopic invariance.* If the symplectic paths σ and σ' are homotopic with fixed endpoints, then $\mu_{Sp}(\sigma, \ell) = \mu_{Sp}(\sigma', \ell)$ for all $\ell \in \Lambda(n)$.
- (S₂) *Additivity by concatenation.* If σ and σ' are two consecutive symplectic paths, and if $\sigma * \sigma'$ is the path σ followed by the path σ' then

$$\mu_{Sp}(\sigma * \sigma', \ell) = \mu_{Sp}(\sigma, \ell) + \mu_{Sp}(\sigma', \ell)$$

for all $\ell \in \Lambda(n)$.

- (S₃) *Zero in the stratum.* If σ and ℓ are such that $\text{Im}(\sigma\ell) \subset \Lambda(n)_\ell$, then $\mu_{Sp}(\sigma, \ell) = 0$.
- (S₄) *Restriction to loops.* If ψ is a loop in Sp , then $\mu_{Sp}(\psi, \ell) = 2m(\psi\ell)$ for all $\ell \in \Lambda(n)$.

As the homotopy axiom (L₁) for the Lagrangian indices that may be replaced by an axiom (L'₁) of homotopy in the strata the axiom (S₁) above may be strengthened by the following statement

- (S'₁) *If the symplectic paths σ and σ' are such that $\sigma\ell$ and $\sigma'\ell$ are homotopic in strata relative to ℓ , then $\mu_{Sp}(\sigma, \ell) = \mu_{Sp}(\sigma', \ell)$.*

The proof of this is identical to that of (L'₁). What is noteworthy is that the data of an intersection index on $\Lambda(n)$ is equivalent to that of an intersection index on $Sp(n)$. Indeed, let μ_Λ be an intersection index on $\Lambda(n)$ and let $\sigma \in \mathcal{C}(Sp(n))$ be a symplectic path. Then the function

$$\mathcal{C}(Sp(n)) \times \Lambda(n) \ni (\sigma, \ell) \mapsto \mu(\sigma\ell, \ell) \in \mathbb{Z} \quad (17)$$

($\sigma\ell$ being the path $t \mapsto \sigma(t)\ell$) is an intersection index on $Sp(n)$.

Conversely, to each intersection index μ_{Sp} we may associate an intersection index $\mu_{\Lambda(n)}$ on $\Lambda(n)$ in the following way. To each $\ell \in \Lambda(n)$ we have a fibration

$$Sp(n) \longrightarrow Sp/St(\ell) = \Lambda(n)$$

($St(\ell)$ being the stabilizer of ℓ in Sp), hence, every path $\lambda \in \mathcal{C}(\Lambda(n))$ can be lifted to a path $\sigma \in \mathcal{C}(Sp(n))$ such that $\lambda = \sigma\ell$. One verifies that the mapping

$$\mu_{\Lambda(n)} : \mathcal{C}(\Lambda(n)) \times \Lambda(n) \ni (\lambda, \ell) \mapsto \mu_{Sp}(\sigma, \ell) \in \mathbb{Z} \quad (18)$$

defines an intersection index on the manifold $\Lambda(n)$.

As in the Lagrangian case, we have a relative uniqueness result for the symplectic path intersection indices

Corollary 12 *Let μ_{Sp} and μ'_{Sp} be two symplectic intersection indices. There exists a function f such that*

$$\mu_{Sp}(\sigma, \ell) = \mu'_{Sp}(\sigma, \ell) + f(\dim(\sigma(0)\ell \cap \ell)) - f(\dim(\sigma(0)\ell \cap \ell)).$$

Proof. It is a direct consequence of the relative uniqueness theorem for Lagrangian intersection indices. ■

4 The indices μ_∞ on $\Lambda_\infty(n)$

We shall in this section use the same notations as in the previous one for paths joining some point ℓ_i to another point ℓ_j in $\Lambda(n)$. Recall that $\Lambda_\infty(n)$ denotes the universal covering of $\Lambda(n)$. We are going to use the intersection index for Lagrangian paths to define an index $\mu_\infty : \Lambda_\infty(n) \times \Lambda_\infty(n) \longrightarrow \mathbb{Z}$.

4.1 The definition of μ_∞

To each Lagrangian intersection index μ_Λ we may associate a 1-cochain μ_∞ on the universal covering $\pi : \Lambda_\infty(n) \longrightarrow \Lambda(n)$ in the following way. We choose once and for all a base point ℓ_0 for $\Lambda(n)$. The elements of $\Lambda_\infty(n)$ are then homotopy classes with fixed endpoints of continuous paths of origin ℓ_0 in $\Lambda(n)$. The projection π from $\Lambda_\infty(n)$ onto $\Lambda(n)$ is defined by $\pi(\ell_\infty) = \ell$ if ℓ_∞ is the homotopy class of a path λ joining ℓ_0 to ℓ .

Definition 13 *The Leray index associated to μ_Λ is the 1-cochain μ_∞ on $\Lambda_\infty(n)$ defined by*

$$\mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = -\mu_{\Lambda(n)}(\lambda_{12}, \ell_0) + \text{sign}(\ell_0, \ell_1, \ell_2) \quad (19)$$

where $\ell_{1,\infty}$ and $\ell_{2,\infty}$ are the homotopy classes of the paths λ_{01} and λ_{02} , respectively, and $\lambda_{12} \sim \lambda_{10} * \lambda_{02}$. Also, sign is the signature cocycle associated to μ_Λ , as defined in Subsection 3.3.

Note that the Leray indices are antisymmetric

$$\mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = -\mu_\infty(\ell_{2,\infty}, \ell_{1,\infty}) \quad (20)$$

for all $(\ell_{1,\infty}, \ell_{2,\infty}) \in \Lambda(n)_\infty^2$ in view of (8) and the fact that sign is antisymmetric.

By theorem 5 and the fact that the definition of the signature cocycle is intrinsic, two Leray indices μ_∞ and μ'_∞ associated to two intersection indices μ_Λ and μ'_Λ are such that

$$\mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) - \mu'_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = f(\dim(\ell_2 \cap \ell_0)) - f(\dim(\ell_1 \cap \ell_0))$$

for some function $f : \mathbb{N} \longrightarrow \mathbb{Z}$.

The Leray indices have the following properties

Proposition 14 (1) *The coboundary of μ_∞ descends to $\Lambda(n)$, i.e. , $\partial\mu_\infty = \pi^* \text{sign}$. That is,*

$$\mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) - \mu_\infty(\ell_{1,\infty}, \ell_{3,\infty}) + \mu_\infty(\ell_{2,\infty}, \ell_{3,\infty}) = \text{sign}(\ell_1, \ell_2, \ell_3) \quad (21)$$

for all $(\ell_{1,\infty}, \ell_{2,\infty}, \ell_{3,\infty}) \in \Lambda_\infty(n)^3$;

(2) The action of $\pi_1(\Lambda(n))$ on μ_∞ is given by

$$\mu_\infty(\gamma_1 \ell_{1,\infty}, \gamma_2 \ell_{2,\infty}) = \mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) + 2(m(\gamma_1) - m(\gamma_2)) \quad (22)$$

for all $\gamma_1, \gamma_2 \in \pi_1(\Lambda(n))$.

Proof. (1) By the definition (19) of μ_∞ we have

$$\begin{aligned} \mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) - \mu_\infty(\ell_{1,\infty}, \ell_{3,\infty}) + \mu_\infty(\ell_{2,\infty}, \ell_{3,\infty}) = \\ -(\mu_\Lambda(\lambda_{12}, \ell_0) - \mu_\Lambda(\lambda_{13}, \ell_0) + \mu_\Lambda(\lambda_{23}, \ell_0)) + \\ \text{sign}(\ell_0, \ell_2, \ell_3) - \text{sign}(\ell_0, \ell_1, \ell_3) + \text{sign}(\ell_0, \ell_2, \ell_3). \end{aligned}$$

Formula (21) follows: by the axioms (L₂) and (L₄), and taking into consideration the fact that $\gamma_3 = \lambda_{31} * \lambda_{12} * \lambda_{23}$ is homotopic to a point, we have

$$\mu_\Lambda(\lambda_{12}, \ell_0) - \mu_\Lambda(\lambda_{13}, \ell_0) + \mu_\Lambda(\lambda_{23}, \ell_0) = 2m(\gamma_3) = 0.$$

By the cocycle property of sign this yields

$$\text{sign}(\ell_0, \ell_2, \ell_3) - \text{sign}(\ell_0, \ell_1, \ell_3) + \text{sign}(\ell_0, \ell_2, \ell_3) = \text{sign}(\ell_1, \ell_2, \ell_3).$$

(2) In view of the antisymmetry of μ_∞ it suffices to show that

$$\mu_\infty(\gamma \ell_{1,\infty}, \ell_{2,\infty}) = \mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) + 2m(\gamma) \quad (23)$$

for all $\gamma \in \pi_1(\Lambda(n))$. Let γ be a loop with origin ℓ_0 . If λ_{01} is a representative of $\ell_{1,\infty}$, then $\gamma * \lambda_{01}$ is a representative of $\gamma \ell_{1,\infty}$ and, by the definition of μ_∞ :

$$\begin{aligned} \mu_\infty(\gamma \ell_{1,\infty}, \ell_{2,\infty}) &= -\mu_\Lambda(\lambda_{10} * \gamma * \lambda_{02}, \ell_0) + \text{sign}(\ell_0, \ell_1, \ell_2) \\ &= -\mu_\Lambda(\lambda_{10} * \lambda_{02}, \ell_0) - 2m(\gamma) + \text{sign}(\ell_0, \ell_1, \ell_2) \\ &= \mu_\infty(\ell_{1,\infty}, \ell_{2,\infty}) + 2m(\gamma). \end{aligned}$$

Formula (23) then follows by the definition of the Maslov index. ■

It is noteworthy that, conversely, the datum of a 1-cochain μ_∞ on $\Lambda_\infty(n)$ satisfying the property (22) above together with a simple topological property, enables us to construct an index, and by theorem 5, *all* the Lagrangian intersection indices (and hence all symplectic intersection indices).

Theorem 15 *Let μ_∞ be a 1-cochain on $\Lambda(n)$ having the property (22), and being locally constant on each of the sets $\Lambda_\infty^2(n; k)$ ($0 \leq k \leq n$) defined in Subsection 1.1. For $(\lambda_{12}, \ell) \in \mathcal{C}(\Lambda(n)) \times \Lambda(n)$ let us define ℓ_∞ , $\ell_{1,\infty}$ and $\ell_{2,\infty}$ in the following way:*

1. ℓ_∞ is an arbitrary element of $\Lambda_\infty(n)$ covering ℓ .
2. $\ell_{1,\infty}$ is the equivalence class of an arbitrary path $\lambda_{01} \in \mathcal{C}(\Lambda(n))$ joining ℓ_0 to ℓ_1 .
3. $\ell_{2,\infty}$ is the equivalence class of $\lambda_{02} = \lambda_{01} * \lambda_{12}$.

Then the formula

$$\mu_\Lambda(\lambda_{12}, \ell) = \mu_\infty(\ell_{2,\infty}, \ell_\infty) - \mu_\infty(\ell_{1,\infty}, \ell_\infty) \quad (24)$$

defines an intersection index. In particular we have $\partial\mu_\infty = \pi^* \text{sign}$.

Proof. That $\mu_{\Lambda(n)}(\lambda_{12}, \ell)$ is independent of the choice of the element ℓ_∞ of the universal covering $\Lambda_\infty(n)$ covering ℓ follows immediately from (22). Indeed, if ℓ'_∞ covers ℓ then there exists an element $\gamma \in \pi_1(\Lambda(n))$ such that $\ell'_\infty = \gamma\ell_\infty$ and hence

$$\begin{cases} \mu_\infty(\ell_{2,\infty}, \ell'_\infty) = \mu_\infty(\ell_{2,\infty}, \ell_\infty) - 2m(\gamma) \\ \mu_\infty(\ell_{1,\infty}, \ell'_\infty) = \mu_\infty(\ell_{1,\infty}, \ell_\infty) - 2m(\gamma) \end{cases}$$

and consequently

$$\mu_\infty(\ell_{2,\infty}, \ell'_\infty) - \mu_\infty(\ell_{1,\infty}, \ell'_\infty) = \mu_\infty(\ell_{2,\infty}, \ell_\infty) - \mu_\infty(\ell_{1,\infty}, \ell_\infty).$$

Let us show that $\mu_\Lambda(\lambda_{12}, \ell)$ is also independent of the choice of λ_{01} and thereby of the choice of the element $\ell_{1,\infty}$ covering ℓ_1 . We replace λ_{01} by some path λ'_{01} and denote $\ell'_{1,\infty}$ the element of $\Lambda_\infty(n)$ it defines. Then, there exists an element γ of $\pi_1(\Lambda(n))$ such that $\ell'_{1,\infty} = \gamma\ell_{1,\infty}$; $\ell_{2,\infty}$ will thus be replaced by $\ell'_{2,\infty} = \gamma\ell_{2,\infty}$. Using once again (22) we have

$$\mu_\infty(\ell'_{2,\infty}, \ell_\infty) - \mu_\infty(\ell'_{1,\infty}, \ell_\infty) = \mu_\infty(\ell_{2,\infty}, \ell_\infty) - \mu_\infty(\ell_{1,\infty}, \ell_\infty).$$

It remains to prove that the function μ_Λ defined by (24) satisfies the axioms (L₁)–(L₄).

- (L₁) Let us replace the path λ_{12} by any path λ'_{12} homotopic (with fixed endpoints) to λ_{12} . Then $\lambda_{02} = \lambda_{01} * \lambda_{12}$ is replaced by a homotopic path $\lambda'_{02} = \lambda_{01} * \lambda'_{12}$ and the homotopy class $\ell_{2,\infty}$ does not change. Consequently, $\mu_\Lambda(\lambda'_{12}, \ell) = \mu_\Lambda(\lambda_{12}, \ell)$.

(L₂) Consider two consecutive paths λ_{12} and λ_{23} . The index $\mu_\Lambda(\lambda_{23}, \ell)$ is given by

$$\mu_\Lambda(\lambda_{23}, \ell) = \mu_\infty(\ell_{3,\infty}, \ell_\infty) - \mu_\infty(\ell'_{2,\infty}, \ell_\infty)$$

where $\ell'_{2,\infty}$ is the homotopy class of an arbitrary path λ'_{02} and $\ell_{3,\infty}$ that of $\lambda'_{02} * \lambda_{23}$. Let us choose $\lambda'_{02} = \lambda_{02}$. Then $\ell'_{2,\infty} = \ell_{2,\infty}$ and we obtain

$$\begin{aligned} \mu_\Lambda(\lambda_{12}, \ell) + \mu_\Lambda(\lambda_{23}, \ell) &= \mu_\infty(\ell_{2,\infty}, \ell_\infty) - \mu_\infty(\ell_{1,\infty}, \ell_\infty) \\ &\quad + \mu_\infty(\ell_{3,\infty}, \ell_\infty) - \mu_\infty(\ell_{2,\infty}, \ell_\infty) \end{aligned}$$

that is,

$$\mu_\Lambda(\lambda_{12}, \ell) + \mu_\Lambda(\lambda_{23}, \ell) = \mu_\Lambda(\lambda_{13}, \ell).$$

(L₃) Let λ_{12} be a path in the strata $\Lambda_\ell(n; k)$ and denote $\ell_\infty(t)$ the equivalence class of $\lambda_{01} * \lambda_{12}(t)$. The mapping $t \mapsto \ell_\infty(t)$ being continuous, the composition mapping $t \mapsto \mu_\infty(\ell_\infty(t), \ell_\infty)$ is locally constant on the interval $[0, 1]$. It follows that it is constant on the same interval since $\Lambda_\ell(n; k)$ is connected. Its value is

$$\mu_\infty(\ell_\infty(0), \ell_\infty) = \mu_\infty(\ell_\infty(1), \ell_\infty)$$

whereof $\mu(\lambda_{12}, \ell) = 0$.

(L₄) Let $\gamma \in \pi_1(\Lambda(n), \ell_0)$. In view of (22) we have the equality

$$\mu_\Lambda(\gamma, \ell) = \mu_\infty(\gamma \ell_{0,\infty}, \ell_\infty) - \mu_\infty(\ell_{0,\infty}, \ell_\infty) = 2m(\gamma).$$

■

The following result allows us to compare the indices $\mu_\Lambda(\lambda, \ell)$ and $\mu_\Lambda(\lambda, \ell')$ corresponding to different choices of Lagrangians ℓ and ℓ' .

Proposition 16 *For all $\lambda_{12} \in \mathcal{C}(\Lambda(n))$ and $(\ell, \ell') \in \Lambda(n)^2$ we have*

$$\mu_\Lambda(\lambda_{12}, \ell) - \mu_\Lambda(\lambda_{12}, \ell') = \text{sign}(\ell_2, \ell, \ell') - \text{sign}(\ell_1, \ell, \ell') \quad (25)$$

Proof. Considering the notations of theorem 15 we have

$$\begin{aligned} \mu_\Lambda(\lambda_{12}, \ell) - \mu_\Lambda(\lambda_{12}, \ell') &= \\ &\mu_\infty(\ell_{2,\infty}, \ell_\infty) - \mu_\infty(\ell_{2,\infty}, \ell'_\infty) - (\mu_\infty(\ell_{1,\infty}, \ell_\infty) - \mu_\infty(\ell_{1,\infty}, \ell'_\infty)) \end{aligned}$$

that is, in view of (21)

$$\begin{aligned} \mu_\Lambda(\lambda_{12}, \ell) - \mu_\Lambda(\lambda_{12}, \ell') &= \\ & - \mu_\infty(\ell_\infty, \ell'_\infty) + \text{sign}(\ell_2, \ell, \ell') - (-\mu_\infty(\ell_\infty, \ell'_\infty) + \text{sign}(\ell_1, \ell, \ell')) \end{aligned}$$

from which (25) immediately follows. ■

4.2 Construction of the canonical index $\bar{\mu}_\infty$

A consequence of the relative uniqueness theorem 5 is that all Lagrangian intersection indices will be known from the moment we have determined one of them, and it follows from theorem 15 that it suffices for that to construct a Leray index.

Using the intersection theory of Lefschetz chains, Leray has constructed in [18] (Ch.I, §2.5) a function

$$m : \Lambda(n)_\infty^2(0) = \{(\ell_{1,\infty}, \ell_{2,\infty}) : \ell_1 \cap \ell_2 = 0\} \longrightarrow \mathbb{Z}$$

such that

$$m(\ell_{1,\infty}, \ell_{2,\infty}) - m(\ell_{1,\infty}, \ell_{3,\infty}) + m(\ell_{2,\infty}, \ell_{3,\infty}) = \text{Inert}(\ell_1, \ell_2, \ell_3) \quad (26)$$

where $\text{Inert}(\ell_1, \ell_2, \ell_3)$ is the index of inertia of the Lagrangian triple (ℓ_1, ℓ_2, ℓ_3) . This index of inertia is defined in the following way. The transversality condition

$$\ell_1 \cap \ell_2 = \ell_2 \cap \ell_3 = \ell_3 \cap \ell_1 = 0$$

being equivalent to

$$\ell_1 \oplus \ell_2 = \ell_2 \oplus \ell_3 = \ell_3 \oplus \ell_1 = X \times X^*$$

the relation $z_1 + z_2 + z_3 = 0$ ($z_1 \in \ell_1$, $z_2 \in \ell_2$, $z_3 \in \ell_3$) defines three quadratic forms

$$z_1 \longmapsto \omega(z_2, z_3), \quad z_2 \longmapsto \omega(z_3, z_1), \quad z_3 \longmapsto \omega(z_1, z_2) \quad (27)$$

such that

$$\omega(z_2, z_3) = \omega(z_3, z_1) = \omega(z_1, z_2).$$

These quadratic forms have the same index of inertia and this index is precisely $\text{Inert}(\ell_1, \ell_2, \ell_3)$.

The function m (which Leray calls “Maslov index”) can be computed using the following formula, due to Souriau [24]:

$$m(\ell_{1,\infty}, \ell_{2,\infty}) = \frac{1}{2\pi} [\theta_1 - \theta_2 + i \text{Tr} \text{Log}(-w_1(w_2^{-1}))] + \frac{n}{2} \quad (28)$$

where $\ell_{1,\infty}$ and $\ell_{2,\infty}$ are identified with (w_1, θ_1) and (w_2, θ_2) , respectively (see 2). Note that

$$\dim(\ell_1 \cap \ell_2) = \text{corang}(w_1 - w_2) \quad (29)$$

(it is the multiplicity of the eigenvalue $+1$ of $w_1(w_2^{-1})$). The logarithm (28) is defined, for every matrix u that has not the eigenvalue -1 , by the formula

$$\text{Log}(u) = \int_{-\infty}^0 \left((\lambda I - u)^{-1} - (\lambda - 1)^{-1} I \right) d\lambda$$

and hence $\text{Log } w_1(w_2^{-1})$ is well-defined since in view of (29)

$$\ell_1 \cap \ell_2 = 0 \iff \det(w_1 - w_2) \neq 0,$$

that is

$$+1 \text{ is not an eigenvalue of } w_1(w_2^{-1}).$$

The function m possesses the following properties:

Proposition 17 (1) *m is locally constant on its domain*

$$\Lambda_{\infty}^2(n; 0) = \{(\ell_{1,\infty}, \ell_{2,\infty}) : \ell_1 \cap \ell_2 = 0\};$$

(2) *The action of $\pi_1(\Lambda(n))$ on m is given by*

$$m(\gamma_1 \ell_{1,\infty}, \gamma_2 \ell_{2,\infty}) = m(\ell_{1,\infty}, \ell_{2,\infty}) + m(\gamma_1) - m(\gamma_2). \quad (30)$$

Proof. (1): Souriaus' formula (28) shows that m is continuous and hence locally constant (it takes its values in a discrete space). (2): To prove the equality (30) we first note that $\pi_1(\Lambda(n)) \equiv \pi_1(W(n))$ can be identified with the group

$$G = \{(I, 2k\pi) : k \in \mathbb{Z}\}$$

acting on $W_{\infty}(n)$ by

$$(I, 2k\pi)(w, \theta) = (w, \theta + 2k\pi).$$

Under this identification, the generator β of $\pi_1(\Lambda(n))$ defined in Proposition 5 is identified with $(I, 2\pi)$. For all $\gamma \in \pi_1(\Lambda(n))$ there exists an integer k such that $\gamma = \beta^k$, and hence

$$\gamma \ell_{\infty} = \beta^k \ell_{\infty} = (w, \theta + 2k\pi) \text{ if } \ell_{\infty} = (w, \theta).$$

In view of the definition (28) of m we thus obtain

$$\begin{aligned} m(\gamma_1 \ell_{1,\infty}, \gamma_2 \ell_{2,\infty}) &= m(\beta^{k_1} \ell_{1,\infty}, \beta^{k_2} \ell_{2,\infty}) - m(\ell_{1,\infty}, \ell_{2,\infty}) \\ &= \frac{1}{2\pi} [2k_1\pi - 2k_2\pi] \\ &= k_1 - k_2 \end{aligned}$$

hence (30) in view of the formulae (4) and (5). ■

Let us now set

$$\tau = 2 \text{Inert} - n. \quad (31)$$

We have the important relation

$$\tau(\ell_1, \ell_2, \ell_3) = \tau^+ - \tau^- \quad (32)$$

where τ^+ (resp. τ^-) is the number of > 0 (resp. < 0) eigenvalues of the quadratic form

$$Q(z_1, z_2, z_3) = \omega(z_1, z_2) + \omega(z_2, z_3) + \omega(z_3, z_1) \quad (33)$$

(see [7]). It follows that τ is the signature of Demazure–Kashiwara (see [6, 19, 20]). It is defined on all triples (ℓ_1, ℓ_2, ℓ_3) , which is not the case for Inert as seen above.

Remark 18 *The function τ is often called “Maslov index”, or “trilateral Maslov index” in the literature (for instance in [4] and [20]). We will not use this terminology.*

The signature of Demazure–Kashiwara has the following properties (see Libermann–Marle [19] or Lion–Vergne [20]; in Capell *et al.* [4] one can find other interesting properties).

Proposition 19 *We have:*

- (1) $\tau \in C_{ant}^2(\Lambda(n), \mathbb{Z})^{Sp}$, i.e. , τ is a 2-cocycle, totally antisymmetric and $Sp(n)$ -invariant :

$$\partial\tau = 0 \quad , \quad \varepsilon^* \tau = (-1)^{sgn(\varepsilon)} \tau \quad , \quad s^* \tau = \tau \quad (34)$$

(ε all permutation of $\{1, 2, 3\}$, $s \in Sp$).

- (2) τ is locally constant on the sets

$$\{(\ell_1, \ell_2, \ell_3) : \dim(\ell_1 \cap \ell_2) = k_1, \dim(\ell_1 \cap \ell_3) = k_2, \dim(\ell_2 \cap \ell_3) = k_3\}$$

- (3) Let A a symmetric matrix of order n and ℓ_A the Lagrangian with equation $p = Ax$. We have

$$\tau(X^*, \ell_A, X) = \text{sign } A \quad (35)$$

where $\text{sign } A$ is the difference between the number of eigenvalues > 0 and < 0 of A .

Let us now set

$$\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = 2m(\ell_{1,\infty}, \ell_{2,\infty}) - n \quad (36)$$

that is, in view of (28) :

$$\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = \frac{1}{\pi} [\theta_1 - \theta_2 + i \operatorname{Tr} \operatorname{Log}(-w_1(w_2^{-1}))]. \quad (37)$$

For $(\ell_{1,\infty}, \ell_{2,\infty}) \in \Lambda_\infty(n)^2$ let us choose $\ell_{3,\infty} \in \Lambda(n)_\infty$ such that

$$\ell_1 \cap \ell_3 = \ell_2 \cap \ell_3 = 0 \quad (38)$$

and *define*

$$\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) = \bar{\mu}_\infty(\ell_{1,\infty}, \ell_{3,\infty}) - \bar{\mu}_\infty(\ell_{2,\infty}, \ell_{3,\infty}) + \tau(\ell_1, \ell_2, \ell_3). \quad (39)$$

We can verify, using the cocycle property of τ that

(1) The right-hand side of (39) does not depend on the choice of $\ell_{3,\infty}$ such that we have (38), which justifies the notation $\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty})$ (see [7] where $\bar{\mu}_\infty$ is denoted by μ).

(2) We have $\partial \bar{\mu}_\infty = \pi^* \tau$, i.e.,

$$\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) - \bar{\mu}_\infty(\ell_{1,\infty}, \ell_{3,\infty}) + \bar{\mu}_\infty(\ell_{2,\infty}, \ell_{3,\infty}) = \tau(\ell_1, \ell_2, \ell_3) \quad (40)$$

for all $(\ell_{1,\infty}, \ell_{2,\infty}, \ell_{3,\infty}) \in \Lambda_\infty(n)^3$.

Definition 20 *We will call $\bar{\mu}_\infty$ the “canonical Leray index” and the Lagrangian and symplectic intersection indices associated to $\bar{\mu}_\infty$ will be call the “canonical intersection indices”. These will be denoted $\bar{\mu}_\Lambda$ and $\bar{\mu}_{Sp}$, respectively.*

The following theorem will show that $\bar{\mu}_\infty$ is indeed a Leray index.

Theorem 21 *The function $\bar{\mu}_\infty$ defined by (39) is the unique 1-cochain on $\Lambda_\infty(n)$ having the following two properties:*

(1) $\partial \bar{\mu}_\infty = \pi^* \tau$.

(2) μ_∞ is locally constant on the sets $\Lambda_\infty^2(n; k)$.

Furthermore, this 1-cochain has the following property:

(3) The action of $\pi_1(\Lambda(n))$ on $\bar{\mu}_\infty$ is given by

$$\bar{\mu}_\infty(\gamma_1 \ell_{1,\infty}, \gamma_2 \ell_{2,\infty}) = \bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) + 2(m(\gamma_1) - m(\gamma_2)). \quad (41)$$

Proof. Let us begin by showing that there exists at most one 1-cochain μ_∞ having the properties (1) and (2). To do so, let us denote by ν the difference $\bar{\mu}_\infty - \bar{\mu}'_\infty$ between two cochains having the stated properties. In this case

$$\nu(\ell_{1,\infty}, \ell_{2,\infty}) = \nu(\ell_{1,\infty}, \ell_{3,\infty}) - \nu(\ell_{2,\infty}, \ell_{3,\infty}).$$

for all $(\ell_{1,\infty}, \ell_{2,\infty}, \ell_{3,\infty})$. We choose ℓ_3 in such a way that $\ell_1 \cap \ell_3 = \ell_2 \cap \ell_3 = 0$. In view of the definition of ν and the axiom (L₃), $\nu(\ell_{1,\infty}, \ell_{2,\infty})$ will remain constant in connected neighborhoods of $\ell_{1,\infty}$ and $\ell_{2,\infty}$. The function ν is thereby locally constant on $\Lambda_\infty(n)^2$. But this space is connected, and having realized this we conclude that ν is in fact constant. Also choosing $\ell_{1,\infty} = \ell_{2,\infty}$ we find that the values of ν is zero, and hence that $\bar{\mu}_\infty = \bar{\mu}'_\infty$.

That μ_∞ possesses property (1) is clear from its construction. Let us therefore concentrate on the proof of property (2). Suppose first that $k = 0$. In view of the Souriau formula (28) and definition (36), the function $\bar{\mu}_\infty$ is continuous on $\Lambda_\infty^2(k)$ and is thereby locally constant since $\bar{\mu}_\infty$ takes discrete values. For arbitrary k , it is enough to choose ℓ_3 transversal to ℓ_1 and ℓ_2 in (39) in such a way that $\bar{\mu}_\infty(\ell_{1,\infty}, \ell_{3,\infty})$ and $\bar{\mu}_\infty(\ell_{2,\infty}, \ell_{3,\infty})$ remain constant, and to apply the property (2) of τ in Proposition 19. Let us finally show that μ_∞ has property (3). Choosing once more ℓ_3 such that $\ell_1 \cap \ell_3 = \ell_2 \cap \ell_3 = 0$, we find that in view of (39) and the fact that for all $\gamma \in \pi_1(\Lambda(n))$ the element $\gamma \ell_\infty$ has the same projection ℓ as ℓ_∞

$$\begin{aligned} \bar{\mu}_\infty(\gamma_1 \ell_{1,\infty}, \gamma_2 \ell_{2,\infty}) &= \bar{\mu}_\infty(\gamma_1 \ell_{1,\infty}, \ell_{3,\infty}) \\ &\quad - \bar{\mu}_\infty(\gamma_2 \ell_{2,\infty}, \ell_{3,\infty}) + \tau(\ell_1, \ell_2, \ell_3). \end{aligned}$$

By (30) and the definition (36) of $\bar{\mu}_\infty$ in the transversal case, we obtain

$$\begin{cases} \bar{\mu}_\infty(\gamma_1 \ell_{1,\infty}, \ell_{3,\infty}) = \bar{\mu}_\infty(\ell_{1,\infty}, \ell_{2,\infty}) + 2m(\gamma_1) \\ \bar{\mu}_\infty(\gamma_2 \ell_{2,\infty}, \ell_{3,\infty}) = \bar{\mu}_\infty(\ell_{2,\infty}, \ell_{3,\infty}) + 2m(\gamma_2) \end{cases}$$

and (41) follows. ■

We point out the fact that Theorem 21 also shows that:

Corollary 22 *The Demazure–Kashiwara index is identical to the signature cocycle, i.e., $\tau = \text{sign}$.*

Proof. This is clear since μ_∞ is a Leray index, so $\partial\mu_\infty = \text{sign}$, and since we also have $\partial\mu_\infty = \tau$. ■

5 Further properties of the intersection indices

We shall see in the following paragraphs that the properties of the canonical Leray index enable us to show several properties of the Lagrangian and symplectic intersection indices. Indeed, we shall for instance see that these intersection indices are symplectic invariant.

5.1 symplectic invariance

As we mentioned the results of the preceding section show that the Lagrangian and symplectic intersection indices $\mu_{\Lambda(n)}$ and μ_{Sp} , respectively, are symplectic invariants. In order to prove this fact we need first to show that the Leray index is invariant under the action of the universal covering space of the symplectic group the following

Lemma 23 *The Leray index is invariant under the action of the universal covering space $Sp_{\infty}(n)$ of $Sp(n)$:*

$$\mu_{\infty}(s_{\infty}\ell_{1,\infty}, s_{\infty}\ell_{2,\infty}) = \mu_{\infty}(\ell_{1,\infty}, \ell_{2,\infty}) \quad (42)$$

for all $(s_{\infty}, \ell_{1,\infty}, \ell_{2,\infty}) \in Sp_{\infty}(n) \times \Lambda(n)_{\infty} \times \Lambda(n)_{\infty}$.

Proof. Consider the mapping

$$\mu'_{\infty} : (\ell_{1,\infty}, \ell_{2,\infty}) \longmapsto \mu_{\infty}(s_{\infty}\ell_{1,\infty}, s_{\infty}\ell_{2,\infty}).$$

Since τ is invariant under the action of $Sp(n)$ we have $\partial\mu'_{\infty} = \tau$. Of course, the function μ'_{∞} is locally constant on $\Lambda_{\infty}^2(n; 0)$ and is therefore identical to μ_{∞} in view of the uniqueness property (1) in Theorem 21. ■

Having shown this we may now state the following essential result. The invariance of the symplectic intersection indices will then follow from their definition.

Theorem 24 *The Lagrangian intersection indices are symplectic invariants, i.e.,*

$$\mu_{\Lambda}(s\lambda_{12}, s\ell) = \mu_{\Lambda}(\lambda_{12}, \ell) \text{ for all } s \in Sp(n). \quad (43)$$

Proof. In view of Theorem 5 and using the fact that

$$\dim(s\ell_1 \cap s\ell_2) = \dim(\ell_1 \cap \ell_2)$$

for all $(s, \ell_1, \ell_2) \in Sp(n) \times \Lambda(n)^2$ it suffices to consider the case in which μ_{Λ} is the canonical intersection index $\bar{\mu}_{\Lambda}$. for $s \in Sp(n)$ we consider an

arbitrary symplectic path $\sigma \in \mathcal{C}(Sp(n))$ joining the identity I of $Sp(n)$ to s . We denote s_∞ the homotopy equivalence class of that path, $s_\infty \in Sp_\infty(n)$. As in Theorem 15 we denote by λ_{01} a path in $\Lambda(n)$ joining ℓ_0 to ℓ_1 and $\ell_{1,\infty}$ its homotopy class. $\ell_{2,\infty}$ is here the homotopy class of the path $\lambda_{01} * \lambda_{12}$. Denoting by λ an arbitrary path joining ℓ_0 to ℓ in $\Lambda(n)$, and ℓ_∞ its homotopy class we find

$$\begin{aligned} s_\infty \ell_{1,\infty} &= \text{class} [t \mapsto \sigma(t) \lambda_{01}(t), 0 \leq t \leq 1] \\ s_\infty \ell_{2,\infty} &= \text{class} \left[t \mapsto \begin{cases} \sigma(t) \lambda_{01}(t), & 0 \leq t \leq \frac{1}{2} \\ \sigma(2t-1) \lambda_{12}(2t-1), & 0 \leq t \leq \frac{1}{2} \end{cases} \right] \\ s_\infty \ell_\infty &= \text{class} [t \mapsto \sigma(t) \lambda(t), 0 \leq t \leq 1]. \end{aligned}$$

Thus, by (24) in Theorem 15 we obtain

$$\bar{\mu}_\infty(s_\infty \ell_{2,\infty}, s_\infty \ell_\infty) - \bar{\mu}_\infty(s_\infty \ell_{1,\infty}, s_\infty \ell_\infty) = \bar{\mu}_\Lambda(s \lambda_{01}, s \ell).$$

But the left-hand side of this equality is

$$\bar{\mu}_\infty(\ell_{2,\infty}, \ell_\infty) - \bar{\mu}_\infty(\ell_{1,\infty}, \ell_\infty) = \bar{\mu}_\Lambda(\lambda_{01}, \ell)$$

by Lemma 23, which proves (43). ■

5.2 The indices $\bar{\mu}_\ell$ on Sp_∞

Theorem 15 expresses every Lagrangian intersection index as the difference between two values of the Leray index on $\Lambda_\infty(n)$. We will now show that there is a similar result for symplectic intersection indices. We begin by showing the following result (*cf.* [7]):

Lemma 25 *Let $s_\infty \in Sp_\infty(n)$ and $\ell_\infty \in \Lambda_\infty(n)$, with projection $\ell \in \Lambda(n)$. The integer $\bar{\mu}_\infty(s_\infty \ell_\infty, \ell_\infty)$ depends solely on (s_∞, ℓ) , and not on the choice of the projection ℓ of ℓ_∞ . Hence, there exists for every $\ell \in \Lambda(n)$, a function $\bar{\mu}_\ell : Sp_\infty(n) \longrightarrow \mathbb{Z}$ such that*

$$\bar{\mu}_\ell(s_\infty) = \bar{\mu}_\infty(s_\infty \ell_\infty, \ell_\infty).$$

Proof. Suppose that ℓ_∞ and ℓ'_∞ are two elements of $\Lambda_\infty(n)$ having the same projection $\ell \in \Lambda(n)$. There is then a $k \in \mathbb{Z}$ such that $\ell'_\infty = \beta^k \ell_\infty$. (β being the generator of $\pi_1(\Lambda(n))$ whose natural image in $\mathbb{Z}$ is $+1$), and in view of (1) we obtain

$$\begin{aligned} \bar{\mu}_\infty(s_\infty \ell'_\infty, \ell'_\infty) &= \bar{\mu}_\infty(s_\infty (\beta^k \ell_\infty), \beta^k \ell_\infty) \\ &= \bar{\mu}_\infty(\beta^k (s_\infty \ell_\infty), \beta^k \ell_\infty) \end{aligned}$$

that is

$$\bar{\mu}_\infty(s_\infty \ell'_\infty, \ell'_\infty) = \bar{\mu}_\infty(s_\infty \ell_\infty, \ell_\infty)$$

by the property (41) in Theorem 21) of $\bar{\mu}_\infty$. ■

We will call the index $\bar{\mu}_\ell$ the “Leray index on $Sp_\infty(n)$ ” relative to the Lagrangian ℓ . We will consider in what follows $s_\infty \in Sp_\infty(n)$ as the homotopy class of a continuous path joining the identity I to an element s (the projection of s_∞) in $Sp(n)$.

Proposition 26 *Let $\sigma_{12} \in \mathcal{C}(Sp(n))$ be a symplectic path joining s_1 to s_2 in $Sp(n)$. Let $s_{1,\infty}$ be an arbitrary element of $Sp_\infty(n)$ covering s_1 and $s_{2,\infty}$ the homotopy class of $\sigma_{01} * \sigma_{12}$ (σ_{01} a representative of $s_{1,\infty}$). We have*

$$\bar{\mu}_{Sp}(\sigma_{12}, \ell) = \bar{\mu}_\ell(s_{2,\infty}) - \bar{\mu}_\ell(s_{1,\infty}). \quad (44)$$

Proof. Consider ℓ_∞ to be the homotopy class of an arbitrary path λ joining ℓ_0 (the base point of $\Lambda(n)_\infty$) to ℓ . We have

$$s_{1,\infty} \ell_\infty = \text{class } [t \mapsto \sigma_{01}(t) \lambda(t), 0 \leq t \leq 1]$$

and

$$s_{2,\infty} \ell_\infty = \text{class } \left[\begin{array}{l} t \mapsto \sigma_{01}(2t) \lambda(2t), 0 \leq t \leq \frac{1}{2} \\ t \mapsto \sigma_{12}(2t-1) \lambda(2t-1), \frac{1}{2} \leq t \leq 1 \end{array} \right]$$

hence, by the formulae (17) and (24) one obtains

$$\begin{aligned} \bar{\mu}_{Sp}(\sigma_{12}, \ell) &= \bar{\mu}_{\Lambda(n)}(\sigma_{12} \ell, \ell) \\ &= \bar{\mu}_\infty(s_{2,\infty} \ell_\infty, \ell_\infty) - \bar{\mu}_\infty(s_{1,\infty} \ell_\infty, \ell_\infty) \end{aligned}$$

that is (44). ■

The two important properties of $\bar{\mu}_{\ell,\infty}$ below are consequences of cohomological property (40) of the Leray index

Proposition 27 *We have*

$$(1) \quad \bar{\mu}_\ell(s_{1,\infty} s_{2,\infty}) = \bar{\mu}_\ell(s_{1,\infty}) + \bar{\mu}_\ell(s_{2,\infty}) + \tau(\ell, s_1 \ell, s_1 s_2 \ell) \quad (45)$$

for all $(s_{1,\infty}, s_{2,\infty}) \in Sp_\infty(n)$.

(2) *Let ℓ and ℓ' be two Lagrangian, then*

$$\bar{\mu}_\ell(s_\infty) - \bar{\mu}_{\ell'}(s_\infty) = \tau(s\ell, \ell, \ell') - \tau(s\ell, s\ell', \ell'). \quad (46)$$

Proof. (1) By the definition of $\bar{\mu}_\ell$ we see that

$$\begin{aligned} \bar{\mu}_\ell(s_{1,\infty}s_{2,\infty}) - \bar{\mu}_\ell(s_{1,\infty}) - \bar{\mu}_\ell(s_{2,\infty}) = \\ \bar{\mu}_\infty(s_{1,\infty}s_{2,\infty}\ell_\infty, \ell_\infty) - \bar{\mu}_\infty(s_{1,\infty}\ell_\infty, \ell_\infty) - \bar{\mu}_\infty(s_{2,\infty}\ell_\infty, \ell_\infty) \end{aligned}$$

that is, using the $Sp_\infty(n)$ -invariance and the antisymmetry of $\bar{\mu}_\infty$:

$$\begin{aligned} \bar{\mu}_\ell(s_{1,\infty}s_{2,\infty}) - \bar{\mu}_\ell(s_{1,\infty}) - \bar{\mu}_\ell(s_{2,\infty}) = \bar{\mu}_\infty(s_{1,\infty}s_{2,\infty}\ell_\infty, \ell_\infty) + \\ \bar{\mu}_\infty(\ell_\infty, s_{1,\infty}\ell_\infty) - \bar{\mu}_\infty(s_{1,\infty}s_{2,\infty}\ell_\infty, s_{1,\infty}\ell_\infty). \end{aligned}$$

In view of the property (40) of $\bar{\mu}_\infty$ we obtain

$$\begin{aligned} \bar{\mu}_\ell(s_{1,\infty}s_{2,\infty}) - \bar{\mu}_\ell(s_{1,\infty}) - \bar{\mu}_\ell(s_{2,\infty}) = \tau(s_1s_2\ell, \ell, s_1\ell) \\ = \tau(\ell, s_1\ell, s_1s_2\ell). \end{aligned}$$

(2) By (40) and the $Sp_\infty(n)$ -invariance of $\bar{\mu}_\infty$,

$$\bar{\mu}_\infty(s_\infty\ell_\infty, \ell_\infty) - \bar{\mu}_\infty(s_\infty\ell_\infty, \ell'_\infty) + \bar{\mu}_\infty(s_\infty\ell_\infty, s_\infty\ell'_\infty) = \tau(s\ell, \ell, \ell')$$

as well as

$$\bar{\mu}_\infty(s_\infty\ell_\infty, s_\infty\ell'_\infty) - \bar{\mu}_\infty(s_\infty\ell_\infty, \ell'_\infty) + \bar{\mu}_\infty(s_\infty\ell'_\infty, \ell'_\infty) = \tau(s\ell, s\ell', \ell')$$

gives us (46) by subtracting the first identity from the second. ■

Corollary 28 *Given two arbitrary Lagrangians ℓ and ℓ' we have*

$$\begin{aligned} \bar{\mu}_{Sp}(\sigma_{12}, \ell) - \bar{\mu}_{Sp}(\sigma_{12}, \ell') &= \tau(s_2\ell, \ell, \ell') - \tau(s_2\ell, s_2\ell', \ell') \\ &\quad - (\tau(s_1\ell, \ell, \ell') - \tau(s_1\ell, s_1\ell', \ell')) \end{aligned} \quad (47)$$

where s_1 and s_2 are the endpoints of the path σ_{12} .

Proof. Formula (44) yields

$$\begin{aligned} \bar{\mu}_{Sp}(\sigma_{12}, \ell) - \bar{\mu}_{Sp}(\sigma_{12}, \ell') = \\ \bar{\mu}_\ell(s_{2,\infty}) - \bar{\mu}_\ell(s_{1,\infty}) - (\bar{\mu}_{\ell'}(s_{2,\infty}) - \bar{\mu}_{\ell'}(s_{1,\infty})) \end{aligned}$$

hence (47), using (46). ■

5.3 Dimensional additivity

The canonical index $\bar{\mu}_\Lambda$ has the interesting “dimensional additivity” property, which is often included as an axiom for the intersection indices (cf. [22]).

Let us introduce the following notations. Let n' and n'' be two integers > 0 and set $n = n' + n''$. We denote by $\Lambda(n')$ and $\Lambda(n'')$ (resp. $Sp(n')$ and $Sp(n'')$) the Lagrangian Grassmannians (resp. the symplectic groups) corresponding to the symplectic spaces $(X' \times X'^*, \omega')$ and $(X'' \times X''^*, \omega'')$. The direct sum

$$\Lambda(n') \oplus \Lambda(n'') = \{\ell' \oplus \ell'' : \ell' \in \Lambda(n'), \ell'' \in \Lambda(n'')\}$$

is identified with a submanifold of $\Lambda(n)$ and

$$Sp(n') \oplus Sp(n'') = \{s' \oplus s'' : s' \in Sp(n'), s'' \in Sp(n'')\}$$

with a subgroup of $Sp(n)$. By definition

$$(s' \oplus s'')(z' \oplus z'') = s'(z') \oplus s''(z''). \quad (48)$$

Denoting τ' and τ'' the signatures of triples of elements of $\Lambda(n')$ and $\Lambda(n'')$ we immediately verify that

$$\tau(\ell'_1 \oplus \ell''_1, \ell'_2 \oplus \ell''_2, \ell'_3 \oplus \ell''_3) = \tau'(\ell'_1, \ell'_2, \ell'_3) + \tau''(\ell''_1, \ell''_2, \ell''_3). \quad (49)$$

In view of the identification of $\Lambda_\infty(n)$ with $W_\infty(n)$ we may define

$$\ell'_\infty \oplus \ell''_\infty = (w' \oplus w'', \theta' + \theta'')$$

if $\ell'_\infty = (w', \theta')$ and $\ell''_\infty = (w'', \theta'')$. This defines an element of $\Lambda(n)_\infty$ since $(w' \oplus w'', \theta' + \theta'') \in W_\infty(n)$.

The following result describes the relation between the Leray indices on the Maslov bundles $\Lambda_\infty(n')$, $\Lambda_\infty(n'')$ and the Leray index on $\Lambda_\infty(n)$

Proposition 29 *Let $\bar{\mu}'_\infty$ and $\bar{\mu}''_\infty$ be the Leray indices on $\Lambda_\infty(n')$ and $\Lambda_\infty(n'')$, respectively. The relation between the Leray indices $\bar{\mu}'_\infty$ and $\bar{\mu}''_\infty$ and the Leray index $\bar{\mu}_\infty$ on $\Lambda_\infty(n)$ is given by*

$$\bar{\mu}_\infty(\ell'_{1,\infty} \oplus \ell''_{1,\infty}, \ell'_{2,\infty} \oplus \ell''_{2,\infty}) = \bar{\mu}'_\infty(\ell'_{1,\infty}, \ell'_{2,\infty}) + \bar{\mu}''_\infty(\ell''_{1,\infty}, \ell''_{2,\infty}). \quad (50)$$

Proof. Let us begin by supposing that $\ell'_{1,\infty}, \ell'_{2,\infty} \in \Lambda_\infty(n')$ and $\ell''_{1,\infty}, \ell''_{2,\infty} \in \Lambda_\infty(n'')$ are transversal: $\ell'_1 \cap \ell'_2 = 0$ and $\ell''_1 \cap \ell''_2 = 0$. Then $\ell'_{1,\infty} \oplus \ell''_{1,\infty}$ and $\ell'_{2,\infty} \oplus \ell''_{2,\infty}$ are also transversal:

$$(\ell'_1 \oplus \ell''_1) \cap (\ell'_2 \oplus \ell''_2) = 0$$

and one shows that

$$\begin{aligned} \text{Log}((w'_1 \oplus w''_1)(w'_2 \oplus w''_2)^{-1}) = \\ \text{Log}(w'_1(w'_2)^{-1}) \oplus \text{Log}(w''_1(w''_2)^{-1}) \end{aligned}$$

under the identifications $\ell'_1 \equiv w'_1$, $\ell'_2 \equiv w'_2$, etc. (see [12]). Formula (50) follows in the considered case. The general case can be reduced to the transversal case by using the definition (39) and the formula (49). ■

From this Proposition we immediately deduce that:

Corollary 30 *Let $\lambda' \in C(\Lambda(n'))$ and $\lambda'' \in C(\Lambda(n''))$ be Lagrangian paths in $\Lambda(n')$ and $\Lambda(n'')$. Then*

$$\lambda' \oplus \lambda'' \in \mathcal{C}(\Lambda(n') \oplus \Lambda(n''))$$

is identified to a Lagrangian path in $\Lambda(n)$ and

$$\bar{\mu}_\Lambda(\lambda' \oplus \lambda'') = \bar{\mu}_{\Lambda'}(\lambda') + \bar{\mu}_{\Lambda''}(\lambda'')$$

where $\bar{\mu}_{\Lambda'}$ and $\bar{\mu}_{\Lambda''}$ are the canonical Lagrangian intersection indices on $\Lambda(n')$ and $\Lambda(n'')$.

Proof. This is trivial by (24) and (50). ■

The case of the symplectic intersection indices is deduced immediately, *mutatis mutandis*:

Corollary 31 *Let $\sigma' \in C(Sp(n'))$ and $\sigma'' \in C(Sp(n''))$ be symplectic paths in $Sp(n')$ and $Sp(n'')$. Then*

$$\sigma' \oplus \sigma'' \in \mathcal{C}(Sp(n') \oplus Sp(n''))$$

is identified with a symplectic path in Sp and

$$\bar{\mu}_{Sp}(\sigma' \oplus \sigma'') = \bar{\mu}_{Sp'}(\sigma') + \bar{\mu}_{Sp''}(\sigma'')$$

where $\bar{\mu}_{Sp'}$ and $\bar{\mu}_{Sp''}$ are the canonical symplectic intersection indices on $Sp(n')$ and $Sp(n'')$.

5.4 The “spectral flow” formula

Let $(A(t))_{0 \leq t \leq 1}$ be a continuous family of real symmetric matrices of order n . We define the “spectral flow” of this continuous family by the formula

$$\text{SF}(A(t))_{0 \leq t \leq 1} = \text{sign } A(1) - \text{sign } A(0). \quad (51)$$

We have the following result, which has been established in a particular case by Duistermaat [13].

Proposition 32 (“Spectral flow formula”) *Let λ_A be the Lagrangian path defined by*

$$\lambda_A(t) = \{x \oplus A(t)x, x \in X\} \quad (52)$$

and σ_A be the symplectic path defined by

$$\sigma_A(t) = \begin{pmatrix} I & 0 \\ A(t) & I \end{pmatrix}$$

for $0 \leq t \leq 1$. Then

$$\text{SF}(A(t))_{0 \leq t \leq 1} = \bar{\mu}_\Lambda(\lambda_A, X) = \bar{\mu}_{Sp}(\sigma_A, X) \quad (53)$$

where $\text{sign } A(t)$ is the difference between the number of eigenvalues $\lambda > 0$ and the number of eigenvalues < 0 of $A(t)$.

Proof. We begin by showing the formula (53). The second one follows immediately since

$$\begin{pmatrix} I & 0 \\ A(t) & I \end{pmatrix} \begin{pmatrix} x \\ 0 \end{pmatrix} = \begin{pmatrix} x \\ A(t)x \end{pmatrix}.$$

Note that $\dim(\lambda_A(t) \cap X^*) = n$ for $0 \leq t \leq 1$, hence

$$\bar{\mu}_\Lambda(\lambda_A, X^*) = 0 \quad (54)$$

in view of the axiom (L_3) which ensures nullity in the strata. In view of formula (25) in Proposition 16 and the antisymmetry of the signature we have

$$\begin{aligned} \bar{\mu}_\Lambda(\lambda_A, X) &= \bar{\mu}_\Lambda(\lambda_A, X) - \bar{\mu}_\Lambda(\lambda_A, X^*) \\ &= \text{sign}(\lambda_A(1), X, X^*) - \text{sign}(\lambda_A(0), X, X^*) \\ &= \text{sign}(X^*, \lambda_A(1), X) - \text{sign}(X^*, \lambda_A(0), X). \end{aligned}$$

Since $\text{sign} = \tau$ we have

$$\bar{\mu}_\Lambda(\lambda_A, X) = \tau(X^*, \lambda_A(0), X) - \tau(X^*, \lambda_A(1), X)$$

and hence

$$\bar{\mu}_\Lambda(\lambda_A, X) = \text{sign } A(1) - \text{sign } A(0)$$

in view of property (35) of τ in Proposition 19. ■

This result is evidently rather trivial in the sense that the spectral flow (51) depends only on the extreme values $A(1)$ and $A(0)$. The situation is however far more complicated in the case of infinite dimensional symplectic spaces and requires elaborated functional analytical techniques (see [3]).

6 Comparative study of some other indices

6.1 The Robbin-Salamon-McDuff index

In [23] (see also [22]) Robbin and Salamon have constructed, using differential geometrical methods, a mapping

$$\mu_{RS} : \mathcal{C}(\Lambda(n)) \times \ell \longrightarrow \frac{1}{2}\mathbb{Z}$$

which they call “Maslov index”. They show that their index possesses the four properties which follow:

RS₁ *If λ and λ' have same endpoints, then $\mu_{RS}(\lambda, \ell) = \mu_{RS}(\lambda', \ell)$ if and only if $\lambda \sim \lambda'$*

RS₂ *$\mu_{RS}(\lambda * \lambda', \ell) = \mu_{RS}(\lambda, \ell) + \mu'_{RS}(\lambda', \ell)$.*

RS₃ *If $\text{Im } \lambda \subset \Lambda_\ell(n; k)$ then $\mu_{RS}(\lambda, \ell) = 0$.*

RS₄ *If γ is a loop in $\Lambda(n)$ then $\mu_{RS}(\gamma, \ell) = m(\gamma)$.*

RS₅ *If λ_A is a Lagrangian path (52), then*

$$\mu_{RS}(\lambda_A, X) = \frac{1}{2} \text{sign } A(1) - \frac{1}{2} \text{sign } A(0). \quad (55)$$

In fact, there is a simple relation between the indices μ_{RS} and $\bar{\mu}_\Lambda$, namely

Proposition 33 *the index μ_{RS} of Robbin-Salamon is given by*

$$\mu_{RS}(\lambda, \ell) = \frac{1}{2} \bar{\mu}_\Lambda(\lambda, \ell)$$

where $\bar{\mu}_\Lambda$ is the canonical intersection index.

Proof. The mapping $2\mu_{RS}$ satisfies the axioms (L₁–L₄) of a Lagrangian intersection index. Therefore, there must exist an index μ_Λ such that $\mu_\Lambda = 2\mu_{RS}$. By Theorem 5 we have thus

$$2\mu_{RS}(\lambda, \ell) = \bar{\mu}_\Lambda(\lambda, \ell) + f(\dim(\lambda(0) \cap \ell)) - f(\dim(\lambda(1) \cap \ell)).$$

for some function f . Taking for λ the path λ_A and $\ell = X$ we have, in view of the formulae (53) and (55) :

$$\begin{aligned} \text{sign } A(1) - \text{sign } A(0) &= \text{sign } A(1) - \text{sign } A(0) + \\ &\quad f(\dim(\lambda_A(0) \cap X)) - f(\dim(\lambda_A(1) \cap X)). \end{aligned}$$

In view of the equality

$$\dim(\lambda_A(t) \cap X) = \text{corang } A(t)$$

this is equivalent to

$$f(\text{corang } A(0)) = f(\text{corang } A(1))$$

for all A , Hence $f = 0$ since $\text{corang } A(t)$ can take arbitrary values. ■

Remark 34 *Robbin and Salamon use in [23] differentiability properties of Lagrangian paths to construct their index. Furthermore, they invoke deep functional analytical properties (Kato's theorem concerning the selection of eigenvalues of a path of symmetric matrices).*

6.2 The Hörmander index

In the frame of his studies of pseudo-differential operators, introduces in [15] a mapping

$$\xi : \Lambda(n)^4 \ni (\ell_1, \ell_2, \ell_3, \ell_4) \longrightarrow \xi(\ell_1, \ell_2, \ell_3, \ell_4) \in \frac{1}{2}\mathbb{Z}.$$

Robbin and Salamon show that the Hörmander index is related to their index μ_{RS} by the formula

$$\xi(\ell_1, \ell_2, \ell_3, \ell_4) = \mu_{RS}(\lambda_{34}, \ell_2) - \mu_{RS}(\lambda_{34}, \ell_1) \quad (56)$$

where λ_{34} is an arbitrary Lagrangian path $\Lambda(n)$ joining ℓ_3 to ℓ_4 .

Proposition 35 *The Hörmander index ξ is given by*

$$\xi(\ell_1, \ell_2, \ell_3, \ell_4) = \frac{1}{2}(\tau(\ell_1, \ell_2, \ell_3) - \tau(\ell_1, \ell_2, \ell_4)) \quad (57)$$

where τ is the Demazure–Kashiwara cocycle.

Proof. In view of Proposition 33 and of (56) we have

$$\xi(\ell_1, \ell_2, \ell_3, \ell_4) = \frac{1}{2}(\bar{\mu}_{\Lambda(n)}(\lambda_{34}, \ell_2) - \bar{\mu}_{\Lambda(n)}(\lambda_{34}, \ell_1)).$$

By the formula (25) in Proposition 16 this is

$$\xi(\ell_1, \ell_2, \ell_3, \ell_4) = \frac{1}{2}(\text{sign}(\ell_4, \ell_2, \ell_1) - \text{sign}(\ell_3, \ell_2, \ell_1))$$

and hence (57), since $\text{sign} = \tau$ and the antisymmetry of τ . ■

References

- [1] Arnold, V. I. *A characteristic class entering in quantization conditions*, Funkt. Anal. i. Priloz. **1**(1), 1–14 (in Russian) (1967); Funct. Anal. Appl. **1**, 1–14 (English translation) (1967).
- [2] Barge, J. and Ghys, E. *Cocycles d'Euler et de Maslov*, Math. Ann. 294, 235–265 (1992).
- [3] Booss-Bavnbek B. and Furutani K. *The Maslov Index: a Functional Analytical Definition and the Spectral Flow Formula*. Tokyo J. Math. **21**(1) (1998).
- [4] Cappell, S. E., Lee R., and Miller, E. Y. *On the Maslov index*, Comm. Pure and Appl. Math. **17** (1994).
- [5] Dazord, P. *Invariants homotopiques attachés aux fibrés symplectiques*, Ann. Inst. Fourier, Grenoble, **29**(2) (1979), 25–78.
- [6] Demazure, M. *Classe de Maslov II*, Exposé numéro 10, Séminaire sur le fibré cotangent, Orsay (1975–76).
- [7] de Gosson, M. *The structure of q-symplectic geometry*, J. Math. Pures et Appl. **71** (1992), 429–453.
- [8] de Gosson, M. *Maslov Classes, Metaplectic Representation and Lagrangian Quantization*, Research Notes in Mathematics **95** (Wiley-VCH, Berlin, 1997).
- [9] de Gosson, M. *Lagrangian path intersections and the Leray index: Aarhus Geometry and Topology Conference*, Contemp. Math. **258** (Amer. Math. Soc., Providence, RI, 2000), 177–184.

- [10] de Gosson, M. and de Gosson, S. *The Cohomological Meaning of Maslov's Lagrangian Path Intersection Index*, Proceedings of the Conference in the Honor of Jean Leray, Karlskrona 1999, Ed. M. de Gosson (Kluwer Acad. Publ., 2001).
- [11] de Gosson, M. and de Gosson, S. *Intersections de chemins dans les variétés de Schubert*. En préparation.
- [12] de Gosson, S. Ph.D. Thesis, Växjö (2006).
- [13] Duistermaat, J. J. *On the Morse index in variational calculus*, Advances in Mathematics **21** (1976), 173–195
- [14] Gutzwiller, M. C. *Chaos in Classical and Quantum Mechanics*, Interdisciplinary Applied Mathematics (Springer-Verlag, 1990).
- [15] Hörmander, L. *Fourier integral operators I*, Acta Mathematica, **127** (1971), 79–183.
- [16] Keller, J. B. *Corrected Bohr-Sommerfeld Quantum Conditions for Non-separable Systems*, Ann. of Physics **4** (1958), 180–188.
- [17] Leray, J. *Complément à la théorie d' Arnold de l'indice de Maslov*, Convegno di geometria симплектика et fisica matematica, Istituto di Alta Matematica, Roma (1973).
- [18] Leray, J. *Lagrangian Analysis and Quantum Mechanics, a mathematical structure related to asymptotic expansions and the Maslov index* (the MIT Press, Cambridge, Mass., 1981); translated from *Analyse Lagrangienne* RCP 25, Strasbourg Collège de France (1976–1977).
- [19] Libermann, P. and Marle, C.-M. *Symplectic Geometry and Analytical Mechanics* (D. Reidel Publishing Company, 1987).
- [20] Lion, G. and Vergne, M. *The Weil representation, Maslov index and Theta series*, Progress in mathematics **6** (Birkhäuser, 1980).
- [21] Maslov, V. P. *Théorie des Perturbations et Méthodes Asymptotiques* (Dunod, Paris, 1972); translated from Russian [original Russian edition: 1965].
- [22] McDuff, D. and Salamon, D. *Symplectic Topology* (Oxford Science Publications, 1998).

- [23] Robbin, J. W. and Salamon, D. A. *The Maslov index for paths*, Topology **32**, 1993, 827–44.
- [24] Souriau, J.-M. *Construction explicite de l'indice de Maslov*, Group Theoretical Methods in Physics, Lecture Notes in Physics, **50** (Springer-Verlag, 1975), 17–148.
- [25] Souriau, J.-M. *Indice de Maslov des variétés lagrangiennes orientables*, C. R. Acad. Sci., Paris, Série A, **276** (1973), 1025–1026.
- [26] Trèves, F. *Introduction to Pseudo-differential and Fourier Integral Operators* (two Volumes), University Series in Mathematics (Plenum Press, 1980).
- [27] Witten, E. in V. I. Arnol'd *et al.* (editors), *Mathematics: Frontiers and Perspectives*, International Mathematical Union, A.M.S (2000), 343.